

How Mycoplasma Spreads

PROJECT TITLE *Mycoplasma bovis* Pneumonia and Arthritis in Feedlot Cattle (6.33)



PROJECT LEADER Dr. Ken Bateman and Dr. Jeff Caswell, Ontario Veterinary College (University of Guelph)

Background

The *Mycoplasma bovis* bacterium has become an important cause of pneumonia and death in Canadian feedlot calves in recent years. *M. bovis* is involved in both the bovine respiratory disease (BRD) complex and the chronic pneumonia and polyarthritis syndrome (CPPS) in high-risk Fall-placed feedlot calves. Exactly how the organism spreads and triggers pneumonia isn't known. This has made it difficult to develop effective prevention and control measures. One challenge has been that some calves carry *M. bovis* in their lungs but do not develop pneumonia, while other infected calves develop a severe and ultimately fatal pneumonia.

Objectives

To determine whether the strain or numbers of *M. bovis* in the lung differ between healthy calves and those with mycoplasma pneumonia.

What they did

A total of 130 newly weaned, non-preconditioned heifer calves were sourced from auction marts and placed in an Ontario feedlot. All calves were given a shot of oxytetracycline and vaccinated against bovine rhinotracheitis, BVD (Types I and II), PI-3 and BRSV at processing. Lung fluid samples were collected from 60 head at arrival. Heifers that developed acute respiratory disease within two weeks of arrival were sampled

again and treated. A healthy control heifer was sampled for each sick heifer sampled. The treated and control heifers were sampled again 60 days after arrival. Lung samples were cultured for *M. bovis*, and bacterial DNA was analyzed to determine the number and type of *M. bovis* bacteria present in sick vs. healthy heifers at each sampling time.

What they learned

Fewer than 2% of calves were infected with *M. bovis* at arrival; this increased to 72% within two weeks, and 86% by 60 days after arrival. This suggests that *M. bovis* infection is uncommon in non-stressed calves prior to weaning, but that the bacteria spread rapidly among calves after they arrive at the feedlot.



How Mycoplasma Spreads continued

Healthy calves were just as likely to be infected with *M. bovis* as calves with pneumonia. The concentration of *M. bovis* bacteria in the lungs of both groups was also similar. This indicates that the presence and number of *M. bovis* bacteria in the lung are not the only factors that determine which calves develop pneumonia.

Minimizing stress, viral infection, and other forms of pneumonia are key to minimizing the occurrence of mycoplasma pneumonia in feedlot calves.

Thirteen strains of *M. bovis* were identified, and the same strains were present in both healthy and sick calves. To verify this, the researchers analyzed *M. bovis* samples from feedlot cattle sampled in a previous BCRC study. As in the present study, the genetic patterns in the *M. bovis* isolated from lungs with pneumonia were similar to those isolated from healthy lungs. Research is ongoing to identify whether virulence differs between the strains.

Several heifers were found to be infected with *M. bovis* at both two weeks and 60 days after arrival, but only one of these heifers was infected by the same strain of *M. bovis* at both time points. This suggests that the calves were able to eliminate one strain of *M. bovis*, but were susceptible to re-infection by a different strain.

What it means

This study improves our understanding of how *M. bovis* enters and spreads within beef feedlots. Calves can successfully clear an *M. bovis* infection without becoming ill. Mycoplasma pneumonia may not develop unless calves are also stressed or infected with other pathogens. The fact that different strains of *M. bovis* have been identified in calves with pneumonia suggests that an effective mycoplasma vaccine will need to provide immunity to multiple strains of *M. bovis*.

These results indicate that minimizing stress, viral infection, and other forms of pneumonia (such as shipping fever) are key to minimizing the occurrence mycoplasma pneumonia in feedlot calves.

Proudly funded by:



The Beef Cattle Research Council (BCRC), a division of the Canadian Cattlemen's Association, sponsors research and technology development and adoption in support of the vision of the Canadian beef industry to have high quality Canadian beef products recognized as the most outstanding by Canadian and world customers.

FOR MORE INFORMATION VISIT:
www.cattle.ca/research

or contact

CANADIAN CATTLEMEN'S ASSOCIATION
#310, 6715 - 8th St. NE, Calgary, AB T2E 7H7
Tel: (403) 275-8558 Fax: (403) 274-5686